

# Genome-wide association mapping of leaf metabolic profiles for dissecting complex traits in maize

Christian Riedelsheimer<sup>a,1</sup>, Jan Lisec<sup>b,1</sup>, Angelika Czedik-Eysenberg<sup>c</sup>, Ronan Sulpice<sup>c,2</sup>, Anna Flis<sup>c</sup>, Christoph Grieder<sup>a</sup>, Thomas Altmann<sup>d</sup>, Mark Stitt<sup>c</sup>, Lothar Willmitzer<sup>b,e</sup>, and Albrecht E. Melchinger<sup>a,3</sup>

<sup>a</sup>Institute of Plant Breeding, Seed Science and Population Genetics, University of Hohenheim, 70593 Stuttgart, Germany; Departments of <sup>b</sup>Molecular Physiology and <sup>c</sup>Metabolic Networks, Max Planck Institute of Molecular Plant Physiology, 14476 Potsdam, Germany; <sup>d</sup>Department Molecular Genetics, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), 06446 Gatersleben, Germany; and <sup>e</sup>King Abdulaziz University, Jeddah 21589, Saudi Arabia

Edited by Edward S. Buckler, US Department of Agriculture, Agriculture Research Service/Cornell University, Ithaca, NY, and accepted by the Editorial Board April 20, 2012 (received for review December 16, 2011)

The diversity of metabolites found in plants is by far greater than in most other organisms. Metabolic profiling techniques, which measure many of these compounds simultaneously, enabled investigating the regulation of metabolic networks and proved to be useful for predicting important agronomic traits. However, little is known about the genetic basis of metabolites in crops such as maize. Here, a set of 289 diverse maize inbred lines was genotyped with 56,110 SNPs and assayed for 118 biochemical compounds in the leaves of young plants, as well as for agronomic traits of mature plants in field trials. Metabolite concentrations had on average a repeatability of 0.73 and showed a correlation pattern that largely reflected their functional grouping. Genome-wide association mapping with correction for population structure and cryptic relatedness identified for 26 distinct metabolites strong associations with SNPs, explaining up to 32.0% of the observed genetic variance. On nine chromosomes, we detected 15 distinct SNP-metabolite associations, each of which explained more than 15% of the genetic variance. For lignin precursors, including *p*-coumaric acid and caffeic acid, we found strong associations (*P* values  $2.7 \times 10^{-10}$  to  $3.9 \times 10^{-18}$ ) with a region on chromosome 9 harboring cinnamoyl-CoA reductase, a key enzyme in monolignol synthesis and a target for improving the quality of lignocellulosic biomass by genetic engineering approaches. Moreover, lignin precursors correlated significantly with lignin content, plant height, and dry matter yield, suggesting that metabolites represent promising connecting links for narrowing the genotype-phenotype gap of complex agronomic traits.

genetic association | metabolomics | *Zea mays*

Plants produce a huge array of biochemical compounds estimated to exceed 200,000 in the plant kingdom (1). Recent progress in analytical capabilities together with advanced data processing techniques enabled the quantitative measurement of hundreds of compounds from a wide range of chemical classes within a single sample of plant material (2). These advances have made it possible to deeply investigate the regulation of metabolic networks and to study their influence on complex traits (3).

Empirical evidence suggested that an array of metabolites can be linked to biomass accumulation in *Arabidopsis thaliana* (4, 5), illustrating their central role for traits connected to growth and development. Metabolomics approaches are also increasingly applied in crop breeding (6). Metabolic profiling could be successfully adopted to predict yield of potato tubers (7) or to distinguish sunflower genotypes with contrasting response to pathogen infections (8). Recently, we showed that metabolic profiles of diverse maize inbred lines allow prediction of their testcross performance in multilocation field trials (9).

Despite these successes, the genetic basis of the metabolic profile in important crops such as maize remains largely unclear. Although certain metabolic products, such as carotenoids in kernels (10), anthocyanins in leaves (11), or maysin (12), have been genetically well characterized, a global picture of the genetic basis of the leaf metabolome is missing. First approaches for studying the genetic basis of concentrations of many distinct metabolites in *Arabidopsis* used populations such as recombinant inbred lines (RIL) that carry

genetic mosaics of two contrasting parental genotypes to map metabolic quantitative trait loci (mQTL) (13). Such linkage mapping approaches revealed a large number of mQTL, but most of them did not explain a substantial amount of genetic variance (14). However, linking genetic variability in metabolite concentrations to genetic variants is of high interest for several reasons.

First, mapping mQTL and ultimately the underlying causal genes can help in annotating the biological function of a metabolite, which may lead to the discovery of new biosynthetic pathways (3). Second, novel enzymatic and regulatory genes controlling metabolic pathways may be identified. Third, mQTL mapping may add functional links to bridge the genotype-phenotype gap of complex traits. In agricultural species, many agronomically important traits are controlled by a large number of genes with small effects (15). Consequently, QTL-based marker-assisted selection is increasingly replaced by whole-genome prediction approaches using thousands of single nucleotide polymorphisms (SNPs) in a black-box prediction model (9, 16). Though this approach is anticipated to be highly successful, it does not provide biological or mechanistic insights into how genetic information is translated into the genetic variability of complex traits. Bridging this apparent genotype-phenotype gap remains a big challenge. A promising approach might therefore be to investigate the genetic basis of intermediate phenotypes with lower genetic complexity, such as yield components or metabolites, and link these results back with the complex trait of interest (17).

Although linkage mapping has a high power for detecting QTL specific to the parental lines of the mapping population, its mapping resolution is very limited due to the few recombination events and, hence, long linkage blocks (18). With the advances in high-throughput genotyping technologies, genome-wide association (GWA) became available as a powerful alternative for dissecting quantitative traits in plants (19). GWA mapping relies on natural linkage disequilibrium (LD) generated by ancestral recombination events in diverse populations. Depending on the level of LD in the population investigated, the mapping resolution can be up to the single nucleotide level. Although plant populations are often prone to inherent population structuring and cryptic relatedness, which can lead to spurious associations in GWA scans (20), powerful techniques are available for decoupling genetic associations with confounding factors (21), and encouraging results have been

Author contributions: T.A., M.S., L.W., and A.E.M. designed research; C.R., J.L., A.C.-E., R.S., A.F., and C.G. performed research; C.R. and J.L. analyzed data; and C.R. and A.E.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. E.S.B. is a guest editor invited by the Editorial Board.


<sup>1</sup>C.R. and J.L. contributed equally to this work.

<sup>2</sup>Present address: Plant Systems Biology Lab, Plant and Agricultural Biosciences Research Centre/Department Botany and Plant Science, National University of Ireland, Galway, Ireland.

<sup>3</sup>To whom correspondence should be addressed. E-mail: melchinger@uni-hohenheim.de.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1120813109/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1120813109/-DCSupplemental).





**Fig. 3.** Manhattan plots for metabolites with known chemical structure and significant association signals. *P* values are shown on a  $\log_{10}$  scale and colored in red if significant with  $\text{FDR} \leq 0.025$ . For the Manhattan plots of the metabolites with unknown chemical structure, see Fig. S1.

containing two sugar/inositol transporter genes as the only genes in this genomic region.

We found three unknown metabolites to show significant associations with an SNP 1.5 kb apart of a cytochrome P450 protein on chromosome 3. Other candidate genes for unknown metabolites included galactinol-sucrose galactosyltransferase, 40S and 60S ribosomal proteins, cellulose synthase-like protein, cysteine synthase, transcription factors, GDSL esterase/lipase, and ubiquitin-associated protein.


The catecholamine dopamine, the phenylpropanoids *p*-coumaric acid, and caffeic acid, as well as two metabolites with unknown chemical structure (1016200-307 and 1044100-307), consistently showed their two strongest significant association signals in a 762-kb region on chromosome 9. To increase the mapping resolution in GWA with a limited amount of SNPs, it has been suggested to impute the allelic states of ungenotyped SNPs based on data from a reference population that has been genotyped at a much higher density (27). We therefore imputed SNPs surrounding the chromosomal region on chromosome 9 using the first-generation HapMap data (1.6 million SNPs) available for 27 maize inbred lines (28). Imputation revealed several closely located SNPs in strong LD and lower *P* values compared with the two surrounding genotyped SNPs located in a cellulose synthase A (*CESA*) and a bZIP transcription factor (*BZIP*) (Fig. 4). For three of the five metabolites (caffeic acid, 1016200-307, and 1044100-307), the strongest signal was consistently observed for a SNP 19.9 kb away from a putative cinnamoyl-CoA reductase (*CCR*), an oxidoreductase important in the monolignol biosynthesis (Fig. 5). This SNP showed also the second lowest *P* value of  $4.1 \times 10^{-12}$  for *p*-coumaric acid. Imputation did not lead to a higher resolution of the other weaker association signals.

*p*-Coumaric acid showed strong negative correlations with dopamine, caffeic acid, 1016200-307, and 1044100-307, which were positively correlated with each other ( $0.65 < r < 0.96$ ; Table 1). These five metabolites with significant signals at the same position on chromosome 9 showed weak but highly significant correlations with the agronomic traits lignin content, plant height, and whole-plant dry matter yield determined in mature

plants grown in the same environment. Correlations with early biomass determined at the time of metabolite measurements were highly significant for four of the five metabolites, but lower compared with the other agronomic traits determined at the end of the vegetation period.

## Discussion

In this study, we showed that GWA mapping is a powerful tool for linking metabolic composition of leaves from field-grown maize inbred lines with genetic variants at a high resolution. Compared with previous mQTL linkage mapping experiments reporting hundreds to several thousands mQTL for the plant metabolome, our results differ concerning both the number of associations and their explained genetic variance. Lisec et al. (13) reported 157 mQTL that account for a median of 4.3% of the phenotypic variation for 181 metabolites with an average repeatability of 0.4. Schauer et al. (29) detected 104 mQTL for



**Fig. 4.** Regional association plot of the region on chromosome 9 for caffeic acid. Imputation revealed several closely located SNPs in strong LD ( $r^2$ ) with the genotyped SNP at position 96,124,914 (black). The strongest association signal was obtained for an SNP 19.9 kb apart from the candidate gene *CCR*.





**Table 1. Correlations between lignin content, plant height, early biomass, dry matter yield, and the five metabolites that show significant associations with the same genomic region on chromosome 9**

	Caffeic acid	Dopamine	<i>p</i> -CA*	1016200-307 <sup>†</sup>	1044100-307 <sup>†</sup>	Lignin	PH	EB	DMY
Caffeic acid	—	0.65	−0.45	0.79	0.72	−0.18	−0.21	−0.18	−0.28
Dopamine	<1 E-15	—	−0.70	0.75	0.69	−0.16	−0.23	−0.13	−0.23
<i>p</i> -CA*	4.4 E-15	<1 E-15	—	−0.72	−0.72	0.15	0.16	0.07	0.12
1016200-307 <sup>†</sup>	<1 E-15	<1 E-15	<1 E-15	—	0.96	−0.20	−0.20	−0.26	−0.33
1044100-307 <sup>†</sup>	<1 E-15	<1 E-15	<1 E-15	<1 E-15	—	−0.19	−0.21	−0.25	−0.35
Lignin ( $w^2 = 0.90$ )	3.3 E-3	6.8 E-3	1.1 E-2	1.0 E-3	1.8 E-3	—	0.30	−0.04	0.07
PH ( $w^2 = 0.96$ )	4.4 E-4	1.6 E-4	8.4 E-3	9.4 E-4	5.5 E-4	6.4 E-7	—	0.00	0.50
EB ( $w^2 = 0.91$ )	3.2 E-3	3.6 E-2	2.8 E-1	3.4 E-5	4.9 E-5	5.3 E-1	9.7 E-1	—	0.45
DMY ( $w^2 = 0.91$ )	2.7 E-6	1.1 E-4	5.0 E-2	2.8 E-8	4.4 E-9	2.4 E-1	<1 E-15	3.1 E-15	—

Pairwise Pearson correlations are shown above the diagonal, and associated *P* values are shown below the diagonal. DMY, dry matter yield; EB, early biomass; PH, plant height.

\**p*-Coumaric acid.

<sup>†</sup>Metabolite with unknown chemical structure.

biofuels through genetic engineering of key regulators in the monolignol synthesis pathway, as suggested by numerous studies (41, 42).

However, the different signs of these correlations illustrate that the relationship between pathway intermediates and lignin content as the final product in the mature plants is not simple and may require consideration of feedback loops. Moreover, the strong correlations of CCR substrates with dopamine, which is known to be stress induced (40), suggests that a change in carbon flux in monolignol synthesis impacts biochemical composition of other secondary metabolites related to, e.g., stress resistance. In fact, the results from several studies showed that perturbing individual steps of the lignin synthesis pathway affects the expression of other genes not only involved in lignin synthesis (43). Although encouraging results of modifying lignin content and composition through down-regulation of CCR have been achieved in poplar (44) and tobacco (45), deeper investigations into the regulatory mechanisms of monolignol biosynthesis seems to be crucial for a successful genetic engineering of lignin synthesis without detrimental side effects on biotic or abiotic stress resistance (46).

Because the generated metabolic profile is a snapshot at a certain moment in time during early development, it would be also of interest to quantitatively measure metabolic fluxes to capture the dynamic component of plant metabolism. Successive measurements of isotope-labeled metabolites have been successfully applied for measuring phenylpropanoids derived from *p*-coumaric acid in potato (47), and a similar approach could shed more light on how lignin synthesis is regulated at the metabolic level in maize.

The established associations with agronomic traits rely on phenotypic correlations of  $|r| \leq 0.35$ , making it difficult to assess quantitatively the direct impact of these genetic variants on the agronomic traits in the field. As expected with a population of our size, the two top significant SNPs on chromosome 9 were not significant in GWA scans of the agronomic traits using a  $Q_{10} + K$  model, and explained less than 1.7% of their genetic variances. Whole-genome prediction, which simultaneously estimates genetic effects over the whole genome instead of focusing on single genomic regions only, remains therefore the method of choice for predicting complex agronomic traits (9).

Given the fact that GWA mapping in elite maize inbred lines is (i) limited in its resolution due to the high level of LD in elite breeding germplasm of maize (9) and (ii) provides only statistical (i.e., indirect) evidence for the association of the genomic region with the investigated metabolites, biological validations of the detected associations remain to be conducted. Possible approaches include RNAi, antisense methods, or the production of knock-out mutants for inducing loss-of-function point mutations in the candidate genes.

In conclusion, we identified strong genetic associations for concentrations of metabolites, especially multiple lignin precursors, to characterize candidate genetic building blocks for lignin content and other agronomic traits. The molecular mechanisms under-

pinning these associations represent promising targets for genetic engineering approaches. Moreover, our results suggest that studying genetically less complex connecting links between genotype and phenotype, such as metabolites, may be a reasonable alternative for GWA mapping of highly complex traits in plants.

## Materials and Methods

**Genetic Material and Field Trials.** The population consisted of the 285 diverse inbred lines described previously (9), with additional four European Flint lines that served as the check genotypes in the field trials. In the trials of each of the three maturity groups, 100 genotypes, including five common check genotypes, were randomized as a  $20 \times 5$   $\alpha$ -lattice design with two replications and planted in two-row plots. Plots were thinned to a final plant density of 100,000 plants per hectare. Early biomass was determined by measuring fresh weight of eight plants per field plot 32 d after sowing. Plant height (m) and dry matter yield of whole-plant biomass (t/ha) were measured for each field plot at the end of the vegetation period. Lignin (%) was measured in the harvested plant material using calibrated near-infrared spectroscopy as described previously (48).

**Metabolic Profiling.** Leaf samples of the inbred lines were collected 33 d after sowing. Samples of ~5 cm were cut from the middle part of the fully developed third leaf of 10 plants per plot, bulked, and immediately frozen using dry ice. The five plots of every incomplete block were sampled within a period of 15 s to minimize within-block error due to metabolic changes over time. All 600 plots were sampled within 69 min. The 50 samples from 10 randomly chosen blocks of one field replication of one maturity group were subsequently processed together as one batch. With this blocking structure, we could account for systematic shifts among batches while keeping the field randomization intact. Analysis of volatizable metabolites was conducted using an established GC-MS method (2) with the assistance of recently developed software (49).

**Genotyping.** Genotyping was performed using the Illumina SNP chip Maize-SNP50 (Illumina Inc.) containing 56,110 unique SNPs. A quality preprocessing was done by applying the following criteria: (i) call rate above 0.95, (ii) unique allele assignment for the 22 replicated checks of genotype B73, (iii) minor allele frequency greater than 2.5%, and (iv) no more than three heterozygous genotypes. A total of 37,227 SNPs met these criteria. Five genotypes with a residual heterozygosity above 5% were excluded. The chromosomal positions of the SNPs refer to the B73 reference genome (B73 RefGen\_v1). Candidate genes were taken from the B73 filtered gene set (release 4a.53).

**Statistical Analysis of Phenotypic and Metabolic Data.** Linear mixed models were used for obtaining least squares means for the phenotypic traits and metabolites. The model for the phenotypic traits was  $y_{ijkl} = \mu + g_i + t_j + r_{jk} + b_{jkl} + e_{ijkl}$ , where  $\mu$  is the grand mean,  $g_i$  the fixed effect of the  $i^{\text{th}}$  genotype,  $t_j$  the fixed effect of the  $j^{\text{th}}$  maturity group trial captured with the common check genotypes,  $r_{jk}$  the fixed effect of the  $k^{\text{th}}$  field replication within the  $j^{\text{th}}$  maturity group trial,  $b_{jkl}$  the random effect of the  $l^{\text{th}}$  incomplete block within the  $jk^{\text{th}}$  field replication, and the residual error  $e_{ijkl} \sim N(0, \sigma_e^2)$ . For the metabolic traits, preprocessing was the same as described previously (9). A fixed effect  $s_{jks}$  for the  $s^{\text{th}}$  batch was included. To

achieve homoscedasticity of the residuals of the metabolites, the flexible Box–Cox power transformation was applied. For each metabolite, the optimum transformation value was determined as described by Piepho (50) using a grid search between 0 and 1 with 100 steps. Repeatabilities ( $w^2$ ) were calculated as  $w^2 = \sigma_g^2 / (\sigma_g^2 + \sigma_e^2 / r)$ , where  $r$  is the number of field replications. Genotypic variance  $\sigma_g^2$  was estimated by restricted maximum likelihood (REML) assuming that  $g_j \sim N(0, \sigma_g^2)$ . REML-based additive estimates of heritabilities were calculated using the function polygenic\_hglm of GenABEL (51) assuming random genotype effects with kinship matrix  $\mathbf{K}$  of proportion of shared SNP alleles as variance-covariance matrix.

**GWA Mapping.** Single-marker analysis was initially carried out using a one-way ANOVA model without considering confounding factors. Phenotypes were regressed on the number of copies of SNP alleles. Quantile–quantile (QQ) plots of the expected vs. observed  $P$  values were inspected for an inflation indicating false positive signals of association. Genome-wide inflation factors ( $\lambda$ ) were calculated as the regression coefficient in the QQ plot with a zero intercept. Because of the high inflation factors, we next applied a  $\mathbf{Q} + \mathbf{K}$  mixed linear model approach with correction for (i) main directions of population structure by regressing on the first three ( $\mathbf{Q}_3$ ) or 10 ( $\mathbf{Q}_{10}$ ) principal components on SNP

data, and (ii) cryptic relatedness using the kinship matrix  $\mathbf{K}$  as variance-covariance matrix for random genotype effects (52). GWA models were fitted using the maximum likelihood implementation in the function polygenic of GenABEL (51).  $P$  values were obtained with the 1 degree of freedom score test implemented in the function mmscore of GenABEL (53).  $P$  values were transformed to  $q$ -values and regarded significant if  $\leq 0.025$  to control for a FDR (25) of 2.5%. The proportion of genetic variance explained by a certain SNP was calculated as  $\rho = R_{LR}^2 / w^2$  using the likelihood-ratio statistic  $R_{LR}^2 = 1 - \exp(-LR/n)$  with  $LR = 2 \times \log(L_{SNP}/L_0)$ , where  $L_0$  is the maximum likelihood of the baseline  $\mathbf{Q}_{10} + \mathbf{K}$  model without considering the SNP, and  $L_{SNP}$  is the maximum likelihood of the full  $\mathbf{Q}_{10} + \mathbf{K}$  model including the SNP as cofactor, and  $n$  is the number of genotypes (54). For the regional association scan on chromosome 9, imputation was performed using BEAGLE 3.3 (55). BEAGLE parameters were set to nSamples = 50 and nIterations = 20.

**ACKNOWLEDGMENTS.** We thank the staff of the experimental research stations of the University of Hohenheim for assistance in conducting the field experiments. This research was supported by the Max Planck Society and the German Federal Ministry of Education and Research (BMBF) within the project GABI-Energy (FKZ: 0315045).

- Dixon RA, Strack D (2003) Phytochemistry meets genome analysis, and beyond. *Phytochemistry* 62:815–816.
- Lisec J, Schauer N, Kopka J, Willmitzer L, Fernie AR (2006) Gas chromatography mass spectrometry-based metabolite profiling in plants. *Nat Protoc* 1:387–396.
- Saito K, Matsuda F (2010) Metabolomics for functional genomics, systems biology, and biotechnology. *Annu Rev Plant Biol* 61:463–489.
- Meyer RC, et al. (2007) The metabolic signature related to high plant growth rate in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 104:4759–4764.
- Sulpice R, et al. (2009) Starch as a major integrator in the regulation of plant growth. *Proc Natl Acad Sci USA* 106:10348–10353.
- Fernie AR, Schauer N (2009) Metabolomics-assisted breeding: A viable option for crop improvement? *Trends Genet* 25:39–48.
- Steinfath M, et al. (2010) Discovering plant metabolic biomarkers for phenotype prediction using an untargeted approach. *Plant Biotechnol J* 8:900–911.
- Peluffo L, et al. (2010) Metabolic profiles of sunflower genotypes with contrasting response to *Sclerotinia sclerotiorum* infection. *Phytochemistry* 71:70–80.
- Riedelsheimer C, et al. (2012) Genomic and metabolic prediction of complex heterotic traits in hybrid maize. *Nat Genet* 44:217–220.
- Wong JC, Lambert RJ, Wurtzel ET, Rocheford TR (2004) QTL and candidate genes phytoene synthase and zeta-carotene desaturase associated with the accumulation of carotenoids in maize. *Theor Appl Genet* 108:349–359.
- Christie PJ, Alfeno MR, Walbot V (1994) Impact of low-temperature stress on general phenylpropanoid and anthocyanin pathways: Enhancement of transcript abundance and anthocyanin pigmentation in maize seedlings. *Planta* 194:541–549.
- Byrne PF, et al. (1996) Quantitative trait loci and metabolic pathways: Genetic control of the concentration of maysin, a corn earworm resistance factor, in maize silks. *Proc Natl Acad Sci USA* 93:8820–8825.
- Lisec J, et al. (2008) Identification of metabolic and biomass QTL in *Arabidopsis thaliana* in a parallel analysis of RIL and IL populations. *Plant J* 53:960–972.
- Brotman Y, et al. (2011) Identification of enzymatic and regulatory genes of plant metabolism through QTL analysis in *Arabidopsis*. *J Plant Physiol* 168:1387–1394.
- Bernardo R (2008) Molecular markers and selection for complex traits in plants: Learning from the last 20 years. *Crop Sci* 48:1649–1664.
- Heffner E, Sorrells M, Jannink JL (2009) Genomic selection for crop improvement. *Crop Sci* 49:1–12.
- Keurentjes JJB (2009) Genetical metabolomics: Closing in on phenotypes. *Curr Opin Plant Biol* 12:223–230.
- Mauricio R (2001) Mapping quantitative trait loci in plants: Uses and caveats for evolutionary biology. *Nat Rev Genet* 2:370–381.
- Stich B, Melchinger A (2010) An introduction to association mapping in plants. *CAB Reviews* 5:1–9.
- Astle W, Balding DJ (2009) Population structure and cryptic relatedness in genetic association studies. *Stat Sci* 24:451–471.
- Sillanpää MJ (2011) Overview of techniques to account for confounding due to population stratification and cryptic relatedness in genomic data association analyses. *Heredity (Edinb)* 106:511–519.
- Atwell S, et al. (2010) Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* 465:627–631.
- Stich B, Melchinger AE (2009) Comparison of mixed-model approaches for association mapping in rapeseed, potato, sugar beet, maize, and *Arabidopsis*. *BMC Genomics* 10:94.
- Aulchenko YS, de Koning DJ, Haley C (2007) Genomewide rapid association using mixed model and regression: A fast and simple method for genomewide pedigree-based quantitative trait loci association analysis. *Genetics* 177:577–585.
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc, B* 57:289–300.
- Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. *Proc Natl Acad Sci USA* 100:9440–9445.
- Marchini J, Howie B (2010) Genotype imputation for genome-wide association studies. *Nat Rev Genet* 11:499–511.
- Gore MA, et al. (2009) A first-generation haplotype map of maize. *Science* 326:1115–1117.
- Schauer N, et al. (2008) Mode of inheritance of primary metabolic traits in tomato. *Plant Cell* 20:509–523.
- Keurentjes JJB, et al. (2006) The genetics of plant metabolism. *Nat Genet* 38:842–849.
- Kim S, et al. (2007) Recombination and linkage disequilibrium in *Arabidopsis thaliana*. *Nat Genet* 39:1151–1155.
- Chan EK, Rowe HC, Kliebenstein DJ (2010) Understanding the evolution of defense metabolites in *Arabidopsis thaliana* using genome-wide association mapping. *Genetics* 185:991–1007.
- Illig T, et al. (2010) A genome-wide perspective of genetic variation in human metabolism. *Nat Genet* 42:137–141.
- Suhre K, et al. (2011) A genome-wide association study of metabolic traits in human urine. *Nat Genet* 43:565–569.
- Fu J, et al. (2009) System-wide molecular evidence for phenotypic buffering in *Arabidopsis*. *Nat Genet* 41:166–167.
- Elshire RJ, et al. (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE* 6:e19379.
- Bonawit ND, Chapple C (2010) The genetics of lignin biosynthesis: Connecting genotype to phenotype. *Annu Rev Genet* 44:337–363.
- Humphreys JM, Chapple C (2002) Rewriting the lignin roadmap. *Curr Opin Plant Biol* 5:224–229.
- Endler A, Persson S (2011) Cellulose synthases and synthesis in *Arabidopsis*. *Mol Plant* 4:199–211.
- Kulma A, Szopa J (2007) Catecholamines are active compounds in plants. *Plant Sci* 172:433–440.
- Chen F, Dixon RA (2007) Lignin modification improves fermentable sugar yields for biofuel production. *Nat Biotechnol* 25:759–761.
- Simmons BA, Loqué D, Ralph J (2010) Advances in modifying lignin for enhanced biofuel production. *Curr Opin Plant Biol* 13:313–320.
- Vanhulme R, Demedts B, Morreel K, Ralph J, Boerjan W (2010) Lignin biosynthesis and structure. *Plant Physiol* 153:895–905.
- Léplé JC, et al. (2007) Downregulation of cinnamoyl-coenzyme A reductase in poplar: Multiple-level phenotyping reveals effects on cell wall polymer metabolism and structure. *Plant Cell* 19:3669–3691.
- Chabannes M, et al. (2001) Strong decrease in lignin content without significant alteration of plant development is induced by simultaneous down-regulation of cinnamoyl CoA reductase (CCR) and cinnamyl alcohol dehydrogenase (CAD) in tobacco plants. *Plant J* 28:257–270.
- Weng JK, Li X, Bonawit ND, Chapple C (2008) Emerging strategies of lignin engineering and degradation for cellulosic biofuel production. *Curr Opin Biotechnol* 19:166–172.
- Matsuda F, et al. (2005) Metabolic flux analysis of the phenylpropanoid pathway in elicitor-treated potato tuber tissue. *Plant Cell Physiol* 46:454–466.
- Grieder C, et al. (2011) Determination of methane fermentation yield and its kinetics by near infrared spectroscopy and chemical composition in maize. *JNIRS* 19:463–477.
- Cuadros-Inostroza A, et al. (2009) TargetSearch—a Bioconductor package for the efficient preprocessing of GC-MS metabolite profiling data. *BMC Bioinformatics* 10:428.
- Piepho H (2009) Data transformation in statistical analysis of field trials with changing treatment variance. *Agron J* 101:865–869.
- Aulchenko YS, Ripke S, Isaacs A, van Duijn CM (2007) GenABEL: An R library for genome-wide association analysis. *Bioinformatics* 23:1294–1296.
- Yu J, et al. (2006) A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nat Genet* 38:203–208.
- Chen WM, Abecasis GR (2007) Family-based association tests for genomewide association scans. *Am J Hum Genet* 81:913–926.
- Sun G, et al. (2010) Variation explained in mixed-model association mapping. *Heredity (Edinb)* 105:333–340.
- Browning BL, Browning SR (2009) A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet* 84:210–223.